



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 639 374 A2**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **94304252.3**

(51) Int. Cl.⁸: **A61K 31/135**

(22) Date of filing: **13.06.94**

(30) Priority: **28.06.93 US 83848**

(43) Date of publication of application:
22.02.95 Bulletin 95/08

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU NL
PT SE**

(71) Applicant: **AMERICAN HOME PRODUCTS
CORPORATION**
Five Giralda Farms
Madison, New Jersey 07940-0874 (US)


(72) Inventor: **Rudolph, Richard Leslie**
978 Heatherstone Drive
Berwyn,
Pennsylvania 19312 (US)
Inventor: **Derivan, Albert Thomas**

226 Hardwicke Lane
Villanova,
Pennsylvania 19085 (US)
Inventor: **Muth, Eric Anthony**
10 Sugaan Close
New Hope,
Pennsylvania 18938 (US)
Inventor: **Upton, Gertrude Virginia**
208 Hermitage Drive
Radnor,
Pennsylvania 19087 (US)

(74) Representative: **Connelly, Michael John et al**
C/o Wyeth Laboratories
Huntercombe Lane South
Taplow
Maldenhead
Berkshire, SL6 0PH (GB)


(54) **New treatments using phenethyl derivatives.**

(57) This invention provides use of a following compound to prepare a medicament for treating obesity, generalized anxiety disorder, post-traumatic stress disorder, late luteal phase dysphoric disorder (premenstrual syndrome), attention deficit disorder, with and without hyperactivity, Gilles de la Tourette syndrome, bulimia nervosa or Shy Drager Syndrome. The compound is a hydroxycycloalkanephenethyl amine of the following structural formula:



in which A is a moiety of the formula

EP 0 639 374 A2



where

the dotted line represents optional unsaturation;

R₁ is hydrogen or alkyl;

R₂ is alkyl;

R₄ is hydrogen, alkyl, formyl, or alkanoyl;

R₅ and R₆ are, independently, hydrogen, hydroxyl, alkyl, alkoxy, alkanoyloxy, cyano, nitro, alkylmercapto, amino, alkylamino, dialkylamino, alkanamido, halo, trifluoromethyl, or taken together, methylene dioxy;

R₇ is hydrogen or alkyl; and

n is 0, 1, 2, 3, or 4;

or a pharmaceutically acceptable salt thereof.

Background of the Invention

The active ingredients of this invention include (1-[2-(dimethylamino)- 1-(4-methoxyphenyl) ethyl] cyclohexanol), or therapeutically acceptable salts thereof, which are known generally as venlafaxine and its analogues. Venlafaxine is disclosed in U. S. Patent No. 4,535,186 (Husbands et al.) and has been previously reported to be useful as an antidepressant. U. S. Patent No. 4,535,186 and US Patent 5,043,460 teach the production of venlafaxine and its analogues and are incorporated herein as reference. For the purposes of this disclosure, and the claims that follow, the use of venlafaxine is understood to include the free base and pharmaceutically acceptable salt forms of venlafaxine, the racemate and its individual enantiomers, and venlafaxine analogs, both as racemates and as their individual enantiomers.

Venlafaxine has been shown to be a potent inhibitor of monoamine neurotransmitter uptake, a mechanism associated with clinical antidepressant activity. Due to its novel structure, venlafaxine has a mechanism of action unrelated to other available antidepressants, such as the tricyclic antidepressants desipramine, nortriptyline, protriptyline, imipramine, amitriptyline, trimipramine and doxepin.

It is believed that venlafaxine's mechanism of action is related to potent inhibition of the uptake of the monoamine neurotransmitters serotonin and norepinephrine. To a lesser degree, venlafaxine also inhibits dopamine reuptake, but it has no inhibitory activity on monoamine oxidase. O-desmethylvenlafaxine, venlafaxine's major metabolite in humans, exhibits a similar pharmacologic profile. Venlafaxine's ability to inhibit norepinephrine and serotonin (5-HT) uptake has been predicted to have an efficacy which rivals or surpasses that of tricyclic antidepressants (Stuart A. Montgomery, M.D., J. Clin. Psychiatry, 54:3, March 1993).

In contrast to classical tricyclic antidepressant drugs, venlafaxine has virtually no affinity for muscarinic, histaminergic or adrenergic receptors in vitro. Pharmacologic activity at these receptors is associated with the various anticholinergic, sedative and cardiovascular effects seen with the tricyclic antidepressant drugs.

Description of the Invention

In accordance with the present invention there is provided use of a compound of the invention (as hereinafter defined) to prepare a medicament for treating, preventing, or controlling obesity, panic disorder, post-traumatic stress disorder, late luteal phase dysphoric disorder (premenstrual syndrome), attention deficit disorders, with and without hyperactivity, Gilles de la Tourette syndrome, bulimia nervosa, generalized anxiety disorder or Shy Drager Syndrome. The medicaments are indicated for use in mammals, preferably in humans. Each of these disorders exhibit a physiological basis for treatment by venlafaxine's ability to inhibit monoamine neurotransmitters.

There is increasing evidence from animal studies that serotonin has an inhibitory role in the control of food intake and modern drug therapy suggests that serotonin is involved in appetite and mood. Both children and adults increase their appetite and gain weight when treated with cyproheptadine, a serotonin receptor blocker. By inhibiting serotonin reuptake, venlafaxine and its analogues can increase synaptic serotonin (5-hydroxytryptamine, 5-HT) and induce anorexia. This action serves as a treatment for obesity.

Bulimia Nervosa is characterized by recurrent binge eating in which the individual feels a loss of control over eating and regularly practices rigorous dieting or fasting, or purging by self-induced vomiting, or the use of diuretics, or laxatives in an attempt to overcome the feeling. Eating binges are generally episodic and may be triggered by psychosocial stress and may occur as often as several times a day. As an antidepressant, venlafaxine can be used to reduce the frequency of bingeing and purging in both depressed and nondepressed bulimics.

Several links exist between late luteal phase dysphoric disorder (LLPDD) and major depression, including similar clinical features and an increased lifetime prevalence of major depression in women with LLPDD. In addition, women with confirmed LLPDD have demonstrated abnormalities in premenstrual levels of serotonin. Premenstrual carbohydrate craving and increased carbohydrate intake in patients with this condition are also suggestive of serotonin involvement. Venlafaxine and its analogues are effective in treating LLPDD because of their serotonin uptake inhibitory ability.

Similarly, the antidepressant activity of venlafaxine and its analogues can be used in the treatment of attention deficit disorders (ADD), with and without hyperactivity, which is characterized by strong behavioral abnormalities. ADD occurs in between 3 and 10% of school age children and is one of the most common childhood and adolescent psychiatric conditions. At the present time, there are three hypotheses suggesting that deficits or dysregulation of the monoamine neurotransmitter system exists in ADD, specifically deficits in noradrenergic, dopaminergic and serotonergic neuronal systems.

Psychopharmacological treatment has proven beneficial in many patients with ADD. The psychotropics most commonly used have come from two different medication groups, the psychostimulants and antidepressants. Imipramine and desipramine have been most commonly prescribed for the treatment of ADD, though fluoxetine has also been used.

One related disorder of interest to the present invention is Gilles de la Tourette syndrome, which is often referred to as Tourette syndrome or Tourette's syndrome. The malady may begin with simple tics, but can progress to multiple, complex movements, which may include vocal and respiratory tics. The vocal tics associated with the syndrome may include grunting or barking noises or may amount to compulsive utterances, often including involuntary curses or derogatory remarks. Agents currently used in the treatment of Tourette syndrome include benzodiazepine anxiolytics, such as lorazepam, for simple tics and haloperidol, clonidine or pimozide for more advanced cases of the syndrome.

Generalized Anxiety Disorder is a syndrome characterized by excessive or chronic anxiety or apprehension concerning two or more of life's circumstances. The disorder's signs and symptoms often include somatic complaints, such as tremor, dyspnea, palpitations, lightheadedness, and nausea.

Acute anxiety attacks (panic disorders) are a defining symptom of anxiety neurosis and are extremely unpleasant for the patient who experiences a subjective fear which arises for no apparent reason. This fear may be a fear of some imminent catastrophe which prevents rational reasoning.


Such anxiety disorders have been treated by a combination of psychologic and pharmacologic measures. Psychologic treatments may include insight psychotherapy, supportive psychotherapy and relaxation techniques, such as meditation or hypnosis. Pharmacologic treatments include those medications that lower the stress level of the patient. Minor tranquilizers are used for controlling the symptoms of chronic or anticipatory anxiety. Panic attacks can be prevented or reduced in severity by therapeutic doses of antidepressants, including venlafaxine and other serotonin reuptake inhibitors, tricyclic antidepressant medications or monoamine oxidase inhibitors.

Venlafaxine and its analogues can also be used to treat Post Traumatic Stress Disorder (PTSD), which may develop after exposure to severe stresses, such as combat, accident, assaults and natural disasters. PTSD is characterized by symptoms of hyperalertness, sleep disturbance, survivor guilt, impairment of concentration and memory, avoidance of reminders and recollection of traumatic events, intensive day-dreams or images, and recurrent nightmares.


While treatment for PTSD consists largely of relaxation techniques designed at relieving the hyperarousal and anxiety symptoms, antidepressants, including serotonin uptake inhibitors, assist in patient recovery.

Venlafaxine and its analogues can also be used as a part of a medical regimen associated with Shy-Drager Syndrome (SDS), which is a multiple systems degeneration resulting in widespread neurologic damage. Autonomic dysfunction with cerebellar ataxia, parkinsonism, corticospinal and corticobulbar tract dysfunction, and amyotrophy are known to occur. The cause of SDS is unknown and its course is progressive. Severe disability or death usually occurs within five to ten years after onset, often from bulbar dysfunction and/or laryngeal stridor.

The use according to the present invention involves preparing a medicament from a compound of the invention. The compounds of invention are substituted phenethylamines. The compounds of this invention are the compounds having the following structural formula:




in which A is a moiety of the formula



where

- 10 the dotted line represents optional unsaturation;
 R_1 is hydrogen or alkyl of 1 to 6 carbon atoms;
 R_2 is alkyl of 1 to 6 carbon atoms;
 R_4 is hydrogen, alkyl of 1 to 6 carbon atoms, formyl, or alkanoyl of 2 to 7 carbon atoms;
 R_5 and R_6 are independently hydrogen, hydroxyl, alkyl of 1 to 6 carbon atoms, alkoxy of 1 to 6
 15 carbon atoms, alkanoyloxy of 2 to 7 carbon atoms, cyano, nitro, alkylmercapto of 1 to 6 carbon atoms, amino, alkylamino of 1 to 6 carbon atoms, dialkylamino in which each alkyl group is of 1 to 6 carbon atoms, alkanamido of 2 to 7 carbon atoms, halo, trifluoromethyl, or when taken together, methylene dioxy;
 R_7 is hydrogen or alkyl of 1 to 6 carbon atoms; and n is 0, 1, 2, 3, or 4;
 20 or a pharmaceutically acceptable salt thereof.

The preferred compounds are those of the formula:



in which

- 35 A is as defined supra;
 R_1 is hydrogen or alkyl of 1 to 3 carbon atoms;
 R_2 is alkyl of 1 to 3 carbon atoms;
 R_5 is hydrogen, hydroxyl, alkoxy of 1 to 3 carbon atoms, chloro, bromo, trifluoromethyl or alkyl of 1
 40 to 3 carbon atoms;
 R_6 is alkyl of 1 to 3 carbon atoms, alkoxy of 1 to 3 carbon atoms, chloro, bromo, trifluoromethyl or alkanoyloxy of 2 to 3 carbon atoms;
 R_7 is hydrogen or alkyl of 1 to 3 carbon atoms;
 or a pharmaceutically acceptable salt thereof.
 45 The most preferred compounds are those in which R_5 and R_6 are in the meta or para positions and n is 2.

Of particular interest are the compounds 1-[(2-dimethylamino)-1-(4-methoxyphenyl)ethyl]cyclohexanol and 1-[(2-dimethylamino)-1-(4-hydroxyphenyl)ethyl]cyclohexanol and the enantiomers and pharmaceutically acceptable salts thereof.

- 50 The compounds in which R_4 is formyl or alkanoyl of 2 to 7 carbon atoms have been found to be not as potent as the corresponding free hydroxy bearing derivatives. However, in long term therapy the acyloxy derivatives will act as pro drugs as the acyl group is removed in vivo either via acid hydrolysis in the stomach or enzymatically.

- The pharmaceutically acceptable acid addition salts of the basic compounds of this invention are
 55 formed conventionally by reaction of the free base with an equivalent amount of any acid which forms a non-toxic salt. Illustrative acids are either inorganic or organic, including hydrochloric, hydrobromic, fumaric, maleic, succinic, sulfuric, phosphoric, tartaric, acetic, citric, oxalic and similar acids. For parenteral administration, the use of water soluble salts is preferred, although either the free base of the pharmaceuti-

cally acceptable salts are applicable for oral or parenteral administration of the antidepressant agents of this invention. The halo substituent representing R_5 or R_6 is intended to include the chloro, bromo, iodo or fluoro substituents.

Pharmaceutical compositions can be prepared from the compounds of this invention. The active ingredient can be compounded into any of the usual oral dosage forms including tablets, capsules and liquid preparations such as elixirs and suspensions containing various coloring, flavoring, stabilizing and flavor masking substances. For compounding oral dosage forms, the active ingredient can be mixed with various conventional tableting materials such as starch, calcium carbonate, lactose, sucrose and dicalcium phosphate to aid the tableting or capsulating process. Magnesium stearate, as an additive, provides a useful lubricant function when desired.

The active ingredients can be dissolved or suspended in a pharmaceutically acceptable sterile liquid carrier, such as sterile water, sterile organic solvent or a mixture of both. Preferably a liquid carrier is one suitable for parenteral injection. Where the active ingredient is sufficiently soluble it can be dissolved in normal saline as a carrier; if it is too insoluble for this it can often be dissolved in a suitable organic solvent, for instance aqueous propylene glycol or polyethylene glycol solutions. Aqueous propylene glycol containing from 10 to 75% of the glycol by weight is generally suitable. In other instances other compositions can be made by dispersing the finely-divided active ingredient in aqueous starch or sodium carboxymethyl cellulose solution, or in a suitable oil, for instance arachis oil. Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by intramuscular, intraperitoneal or subcutaneous injection.

Preferably the pharmaceutical composition is in unit dosage form, e.g. as tablets or capsules. In such form, the composition is sub-divided in unit doses containing appropriate quantities of the active ingredient; the unit dosage forms can be packaged compositions, for example, packeted powders or vials or ampoules. The unit dosage form can be a capsule, cachet or tablet itself, or it can be the appropriate number of any of these in package form. The quantity of the active ingredient in a unit dose of composition may be varied or adjusted from 2 mg. or less to 50 mg. or more, according to the particular need and the activity of the active ingredient. The usual oral recommended dose of venlafaxine for humans may be between about 75 and about 200 mg/day and this dose may be administered in two or three divided doses, preferably with food if administered orally. A maximum recommended daily dose for humans would be about 375 mg, but it will be understood by one skilled in the art that dosage under this invention will be determined by the particular circumstances surrounding each case.

One skilled in this art will also be aware that the routes of administering the compounds of this invention may vary significantly. In addition to other oral administrations, sustained release compositions may be favored. Other acceptable routes may include, but are not limited to, intravenous, intramuscular and intraperitoneal injections, subdermal implants, as well as buccal, sublingual, transdermal, topical, rectal, vaginal and intranasal administrations. Bioerodible, non-bioerodible, biodegradable and non-biodegradable systems of administration may also be used.

It should also be understood that the present invention is intended to include all methods of, and reasons for, treating obesity, panic disorder, post-traumatic stress disorder, late luteal phase dysphoric disorder (premenstrual syndrome), attention deficit disorders, with and without hyperactivity, Gilles de la Tourette syndrome, bulimia nervosa, generalized anxiety disorder or Shy Drager Syndrome in mammals, preferably in humans. For the purposes of this invention, treating these maladies and disorders is to be understood as including all prophylactic, therapeutic, progression inhibiting, remedial, maintenance, curative or other treatments, regimens or administrations of or with venlafaxine that yield the desired effects in the mammal receiving venlafaxine.

The following example is provided to demonstrate the use of venlafaxine in the treatment of obesity. This example is merely illustrative and does not limit the scope of the present invention.

Obesity Reduction Test

A randomized double-blind comparison of venlafaxine and placebo capsules was tested as a treatment for obesity in non-depressed outpatients. The test was conducted over a period of ten (10) weeks with 98 people between the ages of 18 and 65 receiving double-blind medication (50 receiving venlafaxine and 48 receiving a placebo). Each participant in the comparison weighed more than 20%, but less than 100%, above the mean value for sex, height and bone structure according to the 1983 Metropolitan Height and Weight Table. The women in the comparison of childbearing potential received a negative pregnancy test result and agreed to use medically acceptable forms of contraception throughout the period covered by the comparison.

The initial venlafaxine dose was 25 mg at bedtime on study day 1. Subsequently the dose was increased from 50 mg/day to 150 mg/day through study day 14. On study day 15 the dose was increased to 225 mg/day (3 tablets 3 times a day) and this dose was continued for the remainder of the 70 day study. The patients in both groups were also given instructions to decrease their food intake by 20% and to increase their exercise by the same amount. These directives were given to standardize dietary instructions and to try to prevent patients from starting "crash" diets or exercise programs. A specific diet was not prescribed.

Comparisons within groups were performed on changes from baseline weight and body mass index. Body mass index was computed according to the following formula:


$$\text{Body Mass Index} = \frac{\text{Weight (lb.)} \times 100}{\text{Height (in.)} \times \text{Height (in.)}}$$

Obesity Reduction Test Results


The venlafaxine group showed consistent statistically significant mean weight decreases and mean percent decreases from baseline beginning at week 1. Overall, the mean decrease in body weight for the venlafaxine group at week 10 was 7.5 lb with a mean percent decrease from baseline of 3.6%. In contrast, the mean decrease in body weight for the placebo group at week 10 was 1.3 lb with a mean percent decrease from baseline of 0.7%. The body mass index evaluation for the venlafaxine also showed a pattern of decreases similar to that of the weight decreases.

Claims

1. Use of a compound having the formula



in which A is a moiety of the formula



wherein

the dotted line represents optional unsaturation;

R₁ is hydrogen or alkyl of 1 to 6 carbon atoms;

R₂ is alkyl of 1 to 6 carbon atoms;

R₄ is hydrogen, alkyl of 1 to 6 carbon atoms, formyl, or alkanoyl of 2 to 7 carbon atoms;

R₅ and R₆ are independently hydrogen, hydroxyl, alkyl of 1 to 6 carbon atoms, alkoxy of 1 to 6 carbon atoms, alkanoyloxy of 2 to 7 carbon atoms, cyano, nitro, alkylmercapto of 1 to 6 carbon atoms, amino, alkylamino of 1 to 6 carbon atoms, dialkylamino in which

each alkyl group is of 1 to 6 carbon atoms, alkanamido of 2 to 7 carbon atoms, halo, trifluoromethyl, or when taken together, methylene dioxy;

R_7 is hydrogen or alkyl of 1 to 6 carbon atoms; and n is 0, 1, 2, 3, or 4;

or a pharmaceutically acceptable salt thereof, to prepare a medicament for the treatment of obesity, panic disorder, post-traumatic stress disorder, late luteal phase dysphoric disorder, an attention deficit disorder with hyperactivity, an attention deficit disorder without hyperactivity, Gilles de la Tourette syndrome, bulimia nervosa, generalized anxiety disorder, or Shy Drager Syndrome.

2. Use as claimed in claim 1, wherein

R_1 is hydrogen or alkyl of 1 to 3 carbon atoms;

R_2 is alkyl of 1 to 3 carbon atoms;

R_5 is hydrogen, hydroxyl, alkoxy of 1 to 3 carbon atoms, chloro, bromo, trifluoromethyl or alkyl of 1 to 3 carbon atoms;

R_6 is alkyl of 1 to 3 carbon atoms, alkoxy of 1 to 3 carbon atoms, chloro, bromo, trifluoromethyl or alkanoyloxy of 2 to 3 carbon atoms and

R_7 is hydrogen or alkyl of 1 to 3 carbon atoms.

3. Use as claimed in claim 2, wherein R_5 and R_6 are in the meta or para positions and n is 2.

4. Use as claimed in claim 2 wherein the compound is 1-[(2-dimethylamino)-1-(4-methoxyphenyl)ethyl]-cyclohexanol or a pharmaceutically acceptable salt thereof.

5. Use as claimed in claim 2, wherein the compound is 1-[2-(dimethylamino)-1-(4-hydroxyphenyl)ethyl]-cyclohexanol or a pharmaceutically acceptable salt thereof.

6. Use as claimed in any one of claims 1 to 5, wherein the daily dose of the said compound or salt is within the range of 50mg/day to 375mg/day.

7. Use as claimed in claim 4, wherein the daily dose of the said compound or salt is 75mg/day to 200mg/day.



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 639 374 A3**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **94304252.3**

(51) Int. Cl.⁶: **A61K 31/135**

(22) Date of filing: **13.06.94**

(30) Priority: **28.06.93 US 83848**

(43) Date of publication of application:
22.02.95 Bulletin 95/08

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU NL
PT SE**

(88) Date of deferred publication of the search report:
02.08.95 Bulletin 95/31

(71) Applicant: **AMERICAN HOME PRODUCTS
CORPORATION**
Five Giralda Farms
Madison, New Jersey 07940-0874 (US)

(72) Inventor: **Rudolph, Richard Leslie**
978 Heatherstone Drive
Berwyn,
Pennsylvania 19312 (US)

Inventor: **Derivan, Albert Thomas**
226 Hardwicke Lane
Villanova,
Pennsylvania 19085 (US)


Inventor: **Muth, Eric Anthony**
10 Sugaan Close
New Hope,
Pennsylvania 18938 (US)

Inventor: **Upton, Gertrude Virginia**
208 Hermitage Drive
Radnor,
Pennsylvania 19087 (US)

(74) Representative: **Connelly, Michael John et al**
c/o Patent Department
Wyeth Laboratories
Huntercombe Lane South
Taplow
Maidenhead
Berkshire, SL6 0PH (GB)


(54) **New treatments using phenethyl derivatives.**

(57) This invention provides use of a following compound to prepare a medicament for treating obesity, generalized anxiety disorder, post-traumatic stress disorder, late luteal phase dysphoric disorder (premenstrual syndrome), attention deficit disorder, with and without hyperactivity, Gilles de la Tourette syndrome, bulimia nervosa or Shy Drager Syndrome. The compound is a hydroxycycloalkane-phenethyl amine of the following structural formula:



in which A is a moiety of the formula

EP 0 639 374 A3



where

the dotted line represents optional unsaturation;

R_1 is hydrogen or alkyl;

R_2 is alkyl;

R_4 is hydrogen, alkyl, formyl, or alkanoyl;

R_5 and R_6 are, independently, hydrogen, hydroxyl, alkyl, alkoxy, alkanoyloxy, cyano, nitro, alkylmercapto, amino, alkylamino, dialkylamino, alkanamido, halo, trifluoromethyl, or taken together, methylene dioxy;

R_7 is hydrogen or alkyl; and

n is 0, 1, 2, 3, or 4;

or a pharmaceutically acceptable salt thereof.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 94 30 4252

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.5)
X	J. CLIN. PHARMACOL., vol. 32, no.8, 1992 pages 716-724, KAREN J. KLAMERUS ET AL. 'Introduction of a composite parameter to the pharmacokinetics of venlafaxine and its active O-desmethyl metabolite' * page 716, left column, paragraph 1 - right column, paragraph 1 * * abstract *	1-7	A61K31/135
D,Y	EP-A-0 112 669 (AMERICAN HOME PRODUCTS CORPORATION) * page 12, paragraph 3 * * page 14, paragraph 1-3 * * page 16, paragraph 2; claims 1-4 *	1-7	
Y	WO-A-93 09769 (SEPRACOR, INC) * page 1, line 9 - line 25 * * page 3, line 15 - line 17 * * page 5, paragraph 2 *	1-7	
D,Y	J.CLIN.PSYCHIATRY, vol. 54, no.3, 1993 pages 119-126, STUART A. MONTGOMERY 'Venlafaxine: A new dimension in antidepressant pharmacotherapy' * page 119, left column * * page 123, left column, paragraph 3-4 *	1-7	TECHNICAL FIELDS SEARCHED (Int.Cl.5) A61K
Y	DRUGS, vol. 43, no.SUP2, 1992 pages 11-23, S. KASPER ET AL. 'Comparative efficacy of antidepressants' * abstract * * page 17, left column, paragraph 3 *	1-7	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 22 March 1995	Examiner Tzschoppe, D
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons * : member of the same patent family, corresponding document			

EPO FORM 1503 01/92 (P04C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 94 30 4252

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.5)
X	J.CLIN.PSYCHIATRY, vol. 54, no.5SUP, 1993 pages 75-83, STEVEN L. DUBOVSKI 'Approaches to developing new anxiolytics and antidepressants' * page 77, left column, paragraph 3; table 2 * * page 76, right column; table 1 * ---	1-4	
Y	J.CLIN.PSYCHIATRY, vol. 51, no.12supB, 1990 pages 9-12, KARL RICKELS ET AL. 'Clinical overview of serotonin reuptake inhibitors' * the whole document * ---	1-7	
Y	J.CLIN.PSYCHOPHARMACOL., vol. 10, no.2, 1990 pages 152-153, JOHN M. ZAJECKA ET AL. 'Coexisting major depression and obsessive-compulsive disorder treated with venlafaxine' * the whole document * ---	1-7	
Y	AM.J.CLIN.NUTR., vol. 55, no.1sup, 1992 pages 181S-184S, STEPHEN D. WISE 'Clinical studies with fluoxetine in obesity' * abstract * ---	1-7	
Y	DRUGS, vol. 38, no.sup1, 1989 pages 25-31, J.C.PECKOLD ET AL. 'Trimipramine, anxiety,depression and sleep' * abstract * ---	1-7	
			TECHNICAL FIELDS SEARCHED (Int.Cl.5)
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 22 March 1995	Examiner Tzschoppe, D
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure F : intermediate document I : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document			

EPO FORM 1503 (01/92) (P/CA/CI)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 94 30 4252

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.5)
Y	J.CLIN.PSYCHOPHARMACOL., vol. 11, no.4, 1991 pages 233-236, EDWARD SCHWEIZER ET AL. 'Placebo-controlled trial of venlafaxine for the treatment of major depression' * abstract; table 3 * ---	1-7	
Y	EP-A-0 444 855 (ELI LILLY AND COMPANY) 4 September 1991 * page 8, line 3 - line 13 * ---	1-7	
Y	PHARMACOPSYCHIATR., vol. 26, no.suppl, 1993 pages 2-9, B.OLIVIER ET AL. 'Preclinical evidence on the psychotropic profile of fluvoxamine' * abstract * ---	1-7	
Y	J.CLIN.PSYCHIATRY, vol. 52, no.5sup, 1991 pages 52-57, RAY W. FULLER 'Role of serotonin in therapy of depression and related disorders' * abstract * * page 53, left column * --- -/--	1-7	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int.Cl.5)
Place of search MUNICH		Date of completion of the search 22 March 1995	Examiner Tzschoppe, D
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 (04/94) (P0401)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 94 30 4252

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claims	CLASSIFICATION OF THE APPLICATION (Int. Cl. 5)
Y	AM.J.PSYCHIATRY, vol. 47, no.5, 1990 pages 552-564, JAMES I. HUDSON ET AL. 'Affective spectrum disorder: does antidepressant response identify a family of disorders with a common pathophysiology?' * abstract * * page 553, right column, paragraph 5 * * page 558, left column, paragraph 2; table 1 *	1-7	
Y	INT.CLIN.PSYCHOPHARMACOL., vol. 6, no.sup5, 1992 pages 5-12, W.F.BOYER 'Potential indications for the selective serotonin reuptake inhibitors' * abstract *	1-7	
Y	PSYCHOTHER.PSYCHOSOM., vol. 57, no.4, 1992 pages 158-163, MARC DE BOER ET AL. 'Fluvoxamine treatment for chronic PTSD: a pilot study' * abstract *	1-7	TECHNICAL FIELDS SEARCHED (Int. Cl. 5)
Y	J.CLIN.PSYCHOPHARMACOL., vol. 13, no.2, 1993 pages 107-113, LINDA M. NAGY ET AL. 'Open prospective trial of fluoxetine for posttraumatic stress disorder' * page 107 *	1-7	
Y	US-A-5 114 976 (MICHAEL J. NORDEN) 19 May 1992 * claims 1-3 *	1-7	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 22 March 1995	Examiner Tzschoepe, D
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure F : intermediate document T : theory or principle underlying the invention E : earlier patent document, not published on, or after the filing date D : document cited in the application L : document cited for other reasons @ : member of the same patent family, corresponding document			

EPO FORM 1503 (1.9.91) (PUB. NO.)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 94 30 4252

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 5)
Y	WO-A-89 03692 (MASSACHUSETTS INSTITUTE OF TECHNOLOGY) 5 May 1989 * claims 1-17 *	1-7	
X	BR.J.CLIN.PHARMAC., vol. 33, no.6, 1992 pages 589-601, B.SALETU ET AL. 'Pharmacodynamics of venlafaxine evaluated by EEG brain mapping, psychometry and psychophysiology' * table 2 *	1-4	
Y	J.AM.ACAD.CHILD ADLESC.PSYCHIATRY, vol. 32, no.3, May 1993 pages 483-493, PAUL J. AMBROSINI 'Antidepressant treatments in children and adolescents: II. Anxiety, physical, and behavioural disorders' * abstract * * page 485 - page 486 *	1-7	
Y	PEDIATRIC NEUROLOGY, vol. 3, no.3, 1987 pages 129-135, SALLY E. SHAYWITZ 'Attention deficit disorder: current perspectives' * abstract * * page 130 - page 131 *	1-7	
Y	PROG.NEURO-PSYCHOPHARMACOL. & BIOL. PSYCHIAT., vol. 13, no.6, 1989 pages 885-893, MICHAEL J. NORDEN 'Fluoxetine in borderline personality disorder' * abstract * * table 1 *	1-7	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 22 March 1995	Examiner Tzschoppe, D
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 150 (01.94) (PvC01)



European Patent
Office

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claims:
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions, namely:

See sheet -B-

- ☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☒ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respects of which search fees have been paid, namely claims: mentioned in items 1-6,8,9
- ☐ None of the further search fees has been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:



European Patent
Office

EP 94 30 4252 -B-

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions, namely:

1. Claims 1-7 : as far as the treatment of obesity is concerned.
2. Claims 1-7 : as far as the treatment of panic disorder is concerned.
3. Claims 1-7 : as far as the treatment of post-traumatic stress disorder is concerned.
4. Claims 1-7 : as far as the treatment of late luteal phase dysphoric disorder is concerned.
5. Claims 1-7 : as far as the treatment of an attention deficit disorder with hyperactivity is concerned.
6. Claims 1-7 : as far as the treatment of an attention deficit disorder without hyperactivity is concerned.
7. Claims 1-7 : as far as the treatment of Gilles de la Tourette syndrome is concerned.
8. Claims 1-7 : as far as the treatment of bulimia nervosa is concerned.
9. Claims 1-7 : as far as the treatment of generalized anxiety disorder is concerned.
10. Claims 1-7 : as far as the treatment of Shy Drager Syndrome is concerned.

Reasons :

In claim 1 is claimed the use of compounds such as venlafaxine for the manufacture of a medicament for the treatment of 10 different diseases. The common concept linking these different diseases seems to rely in the fact, that all these diseases are treated by inhibition of monoamine neurotransmitters by venlafaxine. As the inhibition of monoamine neurotransmitters (e.g. serotonin reuptake) by venlafaxine and related compounds is already known (see documents of the search report), this common general concept is not novel, and therefore cannot be inventive.